arXiv:1505.00048v1 [math.CT] 30 Apr 2015

PROPs for Linear Systems

Simon Wadsley
Homerton College
Cambridge
CB2 8PH
UK
`S.J.Wadsley@dpmms.cam.ac.uk`

Nick Woods
Department of Mathematics
University of California
Riverside, CA 92521
USA
`woods@math.ucr.edu`

A PROP is a symmetric monoidal category whose objects are the nonnegative integers and whose tensor product on objects is addition. A morphism from m to n in a PROP can be visualized as a string diagram with m input wires and n output wires. For a field k, the PROP $\mathrm{FinVect}_k$ where morphisms are k-linear maps is used by Baez and Erbele to study signal-flow diagrams. We aim to generalize their result characterizing this PROP in terms of generators and relations by looking at the PROP $\mathrm{Mat}(R)$ of matrices with values in R, where R is a commutative rig (that is, a generalization of a ring where the condition that each element has an additive inverse is relaxed). To this end, we show that the category of symmetric monoidal functors out of $\mathrm{Mat}(R)$ is equivalent to the category of bicommutative bimonoids equipped with a certain map of rigs; such functors are called algebras. By choosing R correctly, we will see that the algebras of the PROP FinSpan of finite sets and spans between them are bicommutative bimonoids, while the algebras of the PROP FinRel of finite sets and relations between them are special bicommuative bimonoids and the algebras of $\mathrm{Mat}(\mathbb{Z})$ are bicommutative Hopf monoids.

1 Introduction

Product and permutation categories, or PROPs for short, are tools used to describe the algebraic structure of an object. They were introduced by Mac Lane for the purposes of universal algebra, generalizing Lawvere's notion of an algebraic theory [10, 11].

Formally, a PROP is a symmetric monoidal functor whose objects are the natural numbers and whose tensor product on objects is given by ordinary addition. If $\mathscr{T}$ is a PROP and $\mathscr{C}$ is a symmetric monoidal category, we define an algebra of $\mathscr{T}$ over $\mathscr{C}$ to be a symmetric monoidal functor $\mathscr{T} \to \mathscr{C}$. We say that $\mathscr{T}$ is the PROP for a category $\mathscr{A}$ if the category of algebras over $\mathscr{T}$ is equivalent to $\mathscr{A}$.

PROPs are useful in providing a mathematical formalization of various sorts of diagrams used in physics and engineering, such as signal-flow diagrams [5, 8]. For example, a morphism $m \to n$ can be visualized as a black box diagram with m inputs and n outputs.



The monoidal category structure provides us with different ways of sticking components together — composing two morphisms $f : m \to n$ and $g : n \to p$ lets us connect the outputs of f to the inputs of g, while tensoring provides a way of placing two circuits side-by-side.

Submitted to:
QPL 2015

© Simon Wadsley & Nick Woods
This work is licensed under the
Creative Commons Attribution License.



A rig is generalization of a ring (with unit) where elements are not required to have additive inverses. Some common examples include the natural numbers $\mathbb{N}$ and the two-element Boolean rig $\mathbb{B}$ where 0 and 1 are the truth values FALSE and TRUE, respectively, and where addition is given by OR and multiplication by AND. For a commutative rig R, we define a PROP $\mathrm{Mat}(R)$ by letting morphisms $m \to n$ be $n \times m$ matrices with values in R. Note that $\mathrm{Mat}(R)$ includes some famous categories as particular cases; for example, $\mathrm{Mat}(\mathbb{B})$ is equivalent to the category FinRel of finite sets and relations between them, and $\mathrm{Mat}(\mathbb{N})$ is equivalent to the category FinSpan of finite sets and spans between them. If k is a field, $\mathrm{Mat}(k)$ is equivalent to the category of $\mathrm{FinVect}_k$ of finite k-vector spaces and linear maps between them.

Note that $\mathrm{Mat}(R)$ generalizes Heisenberg's matrix mechanics; $R = \mathbb{C}$ gives the familiar case. If we let $R = [0,\infty)$, then $\mathrm{Mat}(R)$ gives an analogue of matrix mechanics where relative probabilities replace amplitudes. If R is the Boolean rig $\mathbb{B}$, relative possibilities replace amplitudes instead.

We will show:

Theorem. *Let R be a commutative rig. Then* $\mathrm{Mat}(R)$ *is the PROP for bicommutative bimonoids A equipped with a rig map from R to the rig of bimonoid endomorphisms on A.*

In the case that k is the field $\mathbb{R}(s)$ of rational functions in one variable, the string diagrams of the PROP $\mathrm{Mat}(k)$ are the signal-flow diagrams studied by Baez and Erbele and Sobociński et al. [5, 7, 8]. Baez and Erbele show that, when coming up with a set of generators and relations of the category $\mathrm{Mat}(k)$ where k is a field, the additive and multiplicative inverses do not play a role [5]. Instead of describing the PROP in terms of generators and relations, our focus on describing functors out of $\mathrm{Mat}(R)$ give a similar characterization, but the proof is much more efficient. Our method also allows $\mathrm{Mat}(R)$ to be compared to PROPs describing similar structures, such as the one for bimonoids without unit or counit given in [12]. Some consequences of our theorem are that

- FinSpan is the PROP for bicommutative bimonoids;
- FinRel is the PROP for special bicommutative bimonoids; and
- $\mathrm{Mat}(\mathbb{Z})$ is the PROP for bicommutative Hopf monoids.

2 Bicommutative Bimonoids

Fix a monoidal category $\mathscr{C}$ with symmetry functor τ_{AB}. A *commutative monoid* is an object $A \in \mathscr{C}$ with functors $\mu_A : A \otimes A \to A$ and $\eta_A : I \to A$, called multiplication and the unit respectively, satisfying associativity, the unit laws, and commutativity. Diagrammatically, we represent the multiplication and unit operations as



and the relations they must satisfy are depicted as



We define a *monoid homomorphism* to be a function $f : A \to B$ between monoids that commutes with the multiplication and unit maps; that is, $\mu_B(f \otimes f) = f\mu_A$ and $f\eta_A = \eta_B$. This is represented pictorally as



Note that the composition of two monoid homomorphisms is a monoid homomorphism.

A *cocommutative comonoid* $A \in \mathscr{C}$ has the maps comultiplication $\Delta_A : A \to A \otimes A$ and the counit $\varepsilon_A : A \to I$ represented as



These maps satisfy inverted versions of the axioms for a commutative monoid:



We can also define a *comonoid homomorphism*:



Comonoid homomorphisms are again closed under composition.

A *bicommutative bimonoid* A is a commutative monoid and a cocommutative comonoid such that certain distributive laws hold; namely, that maps defining the monoid structure commute with the maps defining the comonoid structure, so that $\mu_A\Delta_A = (\Delta_A \otimes \Delta_A)(1_A \otimes \tau \otimes 1_A)(\mu_A \otimes \mu_A)$, $\Delta_A\eta_A = \eta_A \otimes \eta_A$, $\varepsilon_A\mu_A = \varepsilon_A \otimes \varepsilon_A$, and $\varepsilon_A\mu_A = 1_I$. Pictorally, we have the following relations:



Note that the above axioms are equivalent to saying that Δ_A and ε_A are monoid homomorphisms, or that μ_A and η_A are comonoid homomorphisms. Bimonoid homomorphisms $f : A \to B$ are morphisms that are both monoid and comonoid homomorphisms; they, too, are closed under composition. If $\mathscr{C}$ consists of k-vector spaces and k-linear maps for some field k, then we call the above constructions *commutative algebras*, *cocommutative coalgebras*, and *bicommutative bialgebras* respectively.

Given two bicommutative bimonoids $A,B \in \mathscr{C}$, we can give $A \otimes B$ the structure of a bicommutative bimonoid by setting $\mu_{A\otimes B} = (\mu_A \otimes \mu_B)(1_A \otimes \tau_{BA} \otimes 1_B)$, $\eta_{A\otimes B} = \eta_A \otimes \eta_B$, $\Delta_{A\otimes B} = (1_A \otimes \tau BA \otimes 1_B)(\Delta_A \otimes \Delta_B)$, and $\varepsilon_{A\otimes B} = \varepsilon_A \otimes \varepsilon_B$. In this way we may make $A^{\otimes m}$ into a bicommutative bimonoid for each $m \geq 2$. Since the symmetry τ is natural, the bimonoid homomorphisms of $A \otimes B \to A' \otimes B'$ are exactly $f \otimes g$ where $f : A \to A'$ and $g : B \to B'$ are bimonoid homomorphisms.

We will write μ_A^n for the functor $A^{\otimes n} \to A$ given inductively by $\mu_A^0 = \eta_A$ and $\mu_A^{n+1} = \mu_A(1 \otimes \mu_A^n)$ and Δ_A^n for the functor $A \to A^{\otimes n}$ given inductively by $\Delta_A^0 = \varepsilon_A$ and $\Delta_A^{n+1} = (1_A \otimes \Delta_A^n)\Delta_A$. By associativity, μ_A^n can be depicted as any tree with n leaves; coassociativity means a similar statement holds for Δ_A^n.

We define Bimon to be the subcategory of $\mathscr{C}$ whose objects are bicommutative bimonoids and whose morphisms are the bimonoid homomorphisms between them.

Lemma 1. *The category* Bimon *may be enriched over commutative monoids.*

Proof. If $f,g \in \mathrm{Hom}_{\mathrm{Bimon}}(A,B)$, define $f+g$ to be $\mu_B(f \otimes g)\Delta_A$; note that this is a composition of bimonoid homomorphisms and is hence a bimonoid homomorphism. We need to show that addition defines a commutitive monoid structure on $\mathrm{Hom}_{\mathrm{Bimon}}(A,B)$ and that the composition map $\mathrm{Hom}_{\mathrm{Bimon}}(B,C) \otimes \mathrm{Hom}_{\mathrm{Bimon}}(A,B) \to \mathrm{Hom}_{\mathrm{Bimon}}(A,C)$ is a monoid homomorphism. For the former, associativity occurs since

$$\begin{aligned}(f+g)+h &= \mu_B((\mu_B(f \otimes g)\Delta_A) \otimes h)\Delta_A \\ &= \mu_B(\mu_B \otimes 1)(f \otimes g \otimes h)(\Delta_A \otimes 1)\Delta_A \\ &= \mu_B(1 \otimes \mu_B)(f \otimes g \otimes h)(1 \otimes \Delta_A)\Delta_A \\ &= \mu_B(f \otimes (\mu_B(g \otimes h)\Delta_A)\Delta_A = f+(g+h),\end{aligned}$$

where the third equality follows from the associativity of B and the coassociativity of A. The zero element of $\mathrm{Hom}_{\mathrm{Bimon}}(A,B)$ is $0 = \varepsilon_B\eta_A$; the unit laws of $\mathrm{Hom}_{\mathrm{Bimon}}(A,B)$ follow from the unit laws of B and the counit laws of A, for instance

$$\begin{aligned}0+f &= \mu_B((\varepsilon_B\eta_A) \otimes f)\Delta_A \\ &= \mu_B(\varepsilon_B \otimes 1)f(\eta_A \otimes 1)\Delta_A = f.\end{aligned}$$

To check commutativity, let $\tau_A : A \otimes A \to A \otimes A$ be the symmetry map. Then, by the commutativity of B and the cocommutativity of A,

$$\begin{aligned}f+g &= \mu_B(f \otimes g)\Delta_A = \mu_B\tau_B(f \otimes g)\Delta_A \\ &= \mu_B(g \otimes f)\tau_A\Delta_A = \mu_B(g \otimes f)\Delta_A = g+f.\end{aligned}$$

We now show that the composition map $\mathrm{Hom}_{\mathrm{Bimon}}(B,C) \times \mathrm{Hom}_{\mathrm{Bimon}}(A,B) \to \mathrm{Hom}_{\mathrm{Bimon}}(A,C)$ is bilinear, so it induces a monoid map $\mathrm{Hom}_{\mathrm{Bimon}}(B,C) \otimes \mathrm{Hom}_{\mathrm{Bimon}}(A,B) \to \mathrm{Hom}_{\mathrm{Bimon}}(A,C)$. If $f_1, f_2 \in \mathrm{Hom}_{\mathrm{Bimon}}(B,C)$ and $g_1, g_2 \in \mathrm{Hom}_{\mathrm{Bimon}}(A,B)$, then

$$\begin{aligned}f_1(g_1+g_2) &= f_1\mu_B(g_1 \otimes g_2)\Delta_A \\ &= \mu_C(f_1 \otimes f_1)(g_1 \otimes g_2)\Delta_A \\ &= \mu_C(f_1g_1 \otimes f_1g_2)\Delta_A = f_1g_1 + f_1g_2,\end{aligned}$$

and similarly $(f_1+f_2)g_1 = f_1g_1+f_2g_1$. ☐

Using this enriched structure as addition and composition as multiplication, we can make $\mathrm{Hom}_{\mathrm{Bimon}}(A,A)$ into a rig for all $A \in \mathrm{Bimon}$. We will call this rig $\mathrm{End}(A)$.

For an arbitrary commutative rig R, we define a category Bimon^R whose objects are pairs (A,ϕ_A) where A is a bimonoid in $\mathscr{C}$ and where ϕ_A is a rig homomorphism $R \to \mathrm{End}(A)$. The morphisms $f : (A,\phi_A) \to (B,\phi_B)$ are bimonoid maps $f : A \to B$ for which the diagram

$$\begin{tikzcd}
A \arrow[r, "f"] \arrow[d, "\phi_A(r)"'] & B \arrow[d, "\phi_B(r)"] \\
A \arrow[r, "f"'] & B
\end{tikzcd}$$

commutes for all $r \in R$.

3 The PROP $\mathrm{Mat}(R)$

For a given commutative rig R, we define $\mathrm{Mat}(R)$ to be the PROP where $\mathrm{Hom}_{\mathrm{Mat}(R)}(m,n)$ consists of $n \times m$ R-valued matrices and where composition is defined by matrix multiplication. The tensor product of two matrices $f \otimes g$ is given by the block diagonal matrix

$$\begin{pmatrix} f & 0 \\ 0 & g \end{pmatrix}.$$

Note that, since there is exactly one matrix with n rows and no columns and that its transpose is the unique matrix with no rows and n columns, 0 is both the initial and terminal object in $\mathrm{Mat}(R)$. Also note that the symmetry is given by

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

If f and g are $m \times n$ matrices, we can define an enriched structure on $\mathrm{Mat}(R)$ (see Lemma 1 below) by letting $f+g$ be the usual matrix addition.

We can also equip the object $1 \in \mathrm{Mat}(R)$ with a bicommutative bimonoid structure where (abusing notation slightly) $\mu_R = \begin{pmatrix} 1 & 1 \end{pmatrix}$ and $\eta_R = 0 : 0 \to 1$ are the multiplication and unit maps. The comultiplication Δ_R and counit ε_R are defined by the transposes of these matrices. Checking the axioms now amounts to multiplying matrices; for instance, associativity holds since

$$\mu_R(1 \otimes \mu_R) = \begin{pmatrix} 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \mu_R(\mu_R \otimes 1).$$

Transposing matrices gives a self-inverse contravariant endofunctor on $\mathrm{Mat}(R)$; this functor allows us to extend results shown for the monoid $(\mathrm{Mat}(R),\mu_R,\eta_R)$ to the comonoid $(\mathrm{Mat}(R),\Delta_R,\varepsilon_R)$.

Lemma 2. *For any rig R, the object 1 together with the morphisms μ_R, η_R, Δ_R, and ε_R defined above as well as the map $r : 1 \to 1$ for every $r \in R$ generate* $\mathrm{Mat}(R)$.

Proof. Any object $n \in \mathrm{Mat}(R)$ is equal to $1^{\otimes n}$.

If (r_i) is a $1 \times n$ matrix, then $(r_i) = \mu_R^n((r_1) \otimes \cdots \otimes (r_n))$, since

$$\begin{pmatrix} 1 & \cdots & 1 \end{pmatrix} \begin{pmatrix} r_1 & & 0 \\ & \ddots & \\ 0 & & r_n \end{pmatrix} = \begin{pmatrix} r_1 & \cdots & r_n \end{pmatrix}.$$

If (r_{ij}) is an arbitrary $m \times n$ matrix, we have that $(r_{ij}) = ((r_{i1})_i \otimes \cdots \otimes (r_{in})_i)\Delta_R^m$, where $(r_{ij})_i$ is the jth column of the matrix (r_{ij}), since

$$\begin{pmatrix} (r_{i1})_i & & 0 \\ & \ddots & \\ & & (r_{in})_i \end{pmatrix} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} = \begin{pmatrix} (r_{i1})_i \\ \vdots \\ (r_{in})_i \end{pmatrix}$$

□

4 Algebras over $\mathrm{Mat}(R)$

Lemma 3. *If A is a bicommutative bimonoid, R is a commutative rig, and $\phi : R \to \mathrm{End}(A)$ is a map of rigs, then there is a unique strict monoidal functor F_A enriched over commutative monoids from* Mat(R) *to* Bimon *such that*

(1) $A = F_A(1)$
(2) $\mu_A = F_A(\mu_R)$
(3) $\eta_A = F_A(\eta_R)$
(4) $\Delta_A = F_A(\Delta_R)$
(5) $\varepsilon_A = F_A(\varepsilon_R)$
(6) $\phi(r) = F_A(r : 1 \to 1)$ *for all $r \in R$.*

This functor is necessarily symmetric.

Proof. By Lemma 2 and by the fact that we wish F_A to be a strict monoidal functor, the given information shows how F_A behaves on generators and is therefore enough to define F_A on all of $\mathrm{Mat}(R)$. We are required to set $F_A(1) = A$; on all other objects, the monoidal structre of F_A gives us $F_A(n) = A^{\otimes n}$. Note that F_A is also determined on morphisms; in particular, if (r_i) is a $1 \times n$ matrix, define $F_A(r_i) = \mu_A(\phi(r_1) \otimes \cdots \otimes \phi(r_n))$, and if (r_{ij}) is an arbitrary $m \times n$ matrix, define $F_A(r_{ij}) = (F_A((r_{i1})_i) \otimes \cdots \otimes F_A((r_{im})_i))\Delta^m_{A^{\otimes n}}$. Note that $F_A(r_{ij})$ is a composite of bimonoid maps, and is thus itself a bimonoid map. Note also that, since we have stipulated its values on the generators of $\mathrm{Mat}(R)$, F_A is necessarily unique.

To see that F_A preserves the enriched structure l, first let (r_i) and (s_i) be $1 \times n$ matrices. Then

$$\begin{aligned}
F_A(r_i + s_i) &= \mu_A^n(\phi(r_1 + s_1) \otimes \cdots \otimes \phi(r_n + s_n)) \\
&= \mu_A^n((\phi(r_1) + \phi(s_1)) \otimes \cdots \otimes (\phi(r_n) + \phi(s_n))) \\
&= \mu_A^n(\mu_A(\phi(r_1) \otimes \phi(s_1))\Delta_A \otimes \cdots \otimes \mu_A(\phi(r_n) \otimes \phi(s_n))\Delta_A) \\
&= \mu_A^n \mu_{A^{\otimes n}}(\phi(r_1) \otimes \phi(s_1) \otimes \cdots \otimes \phi(r_n) \otimes \phi(s_n))\Delta_A \\
&= \mu_A^n(\mu_A^n(\phi(r_1) \otimes \cdots \otimes \phi(r_n)) \otimes \mu_A^n(\phi(s_1) \otimes \cdots \otimes \phi(s_n)))\Delta_{A^{\otimes n}} \\
&= F_A(r_i) + F_A(s_i).
\end{aligned}$$

For the general case, assume (r_{ij}) and (s_{ij}) are $m\times n$ matrices. Then

$$\begin{aligned}
F_A(r_{ij}+s_{ij}) &= (F_A(r_{i1}+s_{i1})\otimes\cdots\otimes F_A(r_{im}+s_{im}))\Delta^m_{A^{\otimes n}}\\
&= ((F_A(r_{i1})+F_A(s_{i1}))\otimes\cdots\otimes(F_A(r_{im}+s_{im})))\Delta^m_{A^{\otimes n}}\\
&= (\mu_{A^{\otimes n}}(F_A(r_{i1})\otimes F_A(s_{i1}))\Delta_{A^{\otimes n}}\otimes\cdots\otimes\mu_{A^{\otimes n}}(F_A(r_{i1})\otimes F_A(s_{i1}))\Delta_{A^{\otimes n}})\Delta^m_{A^{\otimes n}}\\
&= \mu_{A^{\otimes m}}((F_A(r_{i1})\otimes F_A(r_{im}))\Delta^m_{A^{\otimes n}}\otimes(F_A(s_{i1})\otimes F_A(s_{im}))\Delta^m_{A^{\otimes n}})\Delta_{A^{\otimes n}}\\
&= \mu_{A^{\otimes m}}(F_A(r_{ij})\otimes F_A(s_{ij}))\Delta_{A^{\otimes n}}\\
&= F_A(r_{ij})+F_A(s_{ij}).
\end{aligned}$$

We now show F_A is a functor. Because we already checked that F_A preserves the enriched structure, it suffices to show that F_A preserves composition for matrices of the form

$$E^{mn}_{ij}(r) = 0^{i-1\times j-1}\otimes(r)\otimes 0^{m-i\times n-j},$$

since every matrix is the sum of matrices which are zero everywhere except at one place. Note that $F_A(E^{1n}_{1j}(r)) = \varepsilon_{A^{\otimes j-1}}\otimes\phi(r)\otimes\varepsilon_{A^{\otimes n-j}}$, so

$$F_A(E^{mn}_{ij}(r)) = \eta_{A^{\otimes j-1}}\otimes\varepsilon_{A^{\otimes i-1}}\otimes\phi(r)\otimes\varepsilon_{A^{\otimes n-i}}\otimes\eta_{A^{\otimes m-j}}.$$

To see that F_A is symmetric, we have, by the naturality of τ_{AA},

$$\begin{aligned}
F_A(E^{22}_{12}(1)+E^{22}_{21}(1)) &= (\eta_A\otimes 1_A\otimes\varepsilon_A)+(\varepsilon_A\otimes 1_A\otimes\eta_A)\\
&= \mu_{A\otimes A}(\eta_A\otimes 1_A\otimes\varepsilon_A\otimes\varepsilon_A\otimes 1_A\otimes\eta_A)\Delta_{A\otimes A}\\
&= (\mu_A\otimes\mu_A)(1_A\otimes\tau_{AA}\otimes 1_A)(\eta_A\otimes 1_A\otimes\varepsilon_A\otimes\varepsilon_A\otimes 1_A\otimes\eta_A)(1_A\otimes\tau_{AA}\otimes 1_A)(\Delta_A\otimes\Delta_A)\\
&= (\mu_A\otimes\mu_A)(\eta_A\otimes 1_A\otimes 1_A\otimes\eta_A)\tau_{AA}(1_A\otimes\eta_A\otimes\eta_A\otimes 1_A)(\Delta_A\otimes\Delta_A)\\
&= (\mu_A(\eta_A\otimes 1_A)\otimes\mu_A(1_A\otimes\eta_A))\tau_{AA}((1_A\otimes\varepsilon_A)\Delta_A\otimes(\varepsilon_A\otimes 1_A)\Delta_A)\\
&= \tau_A.
\end{aligned}$$

□

Lemma 4. *If R is a commutative rig and $F:\mathrm{Mat}(R)\to\mathscr{C}$ is an algebra over the PROP $\mathrm{Mat}(R)$, then there exists some $(A,\phi_A)\in\mathrm{Bimon}^R$ such that $A=F(1)$, $\mu_A=F(\mu_R)$, $\eta_A=F(\eta_R)$, $\Delta_A=F(\Delta_R)$, $\varepsilon_A=F(\varepsilon_R)$, and $\phi_A(r)=F(r:1\to 1)$. Furthermore, F is an enriched functor.*

Proof. The axioms for A being a bicommutative bimonoid follow from the respective axioms of $\mathrm{Mat}(R)$ and the fact that F is a monoidal functor.

We still need to show that F is an enriched functor. Let X and Y be $m\times n$ matrices; then

$$\begin{aligned}
\mu_{A^{\otimes m}}(F(X)\otimes F(Y))\Delta_{A^{\otimes n}} &= F\begin{pmatrix} I_m & I_m\end{pmatrix} F\begin{pmatrix} X & 0\\ 0 & Y\end{pmatrix} F\begin{pmatrix} I_n\\ I_n\end{pmatrix}\\
&= F(X+Y).
\end{aligned}$$

□

We now prove our main theorem:

Theorem 5. *If R is a commutative rig, then $\mathrm{Mat}(R)$ is the PROP for bicommutative bimonoid A equipped with a map of rigs $\phi:R\to\mathrm{End}(A)$.*

Proof. We begin by giving a functor $\mathbf{F}$ from Bimon^R to the category of algebras over $\mathrm{Mat}(R)$. Given an element $(A,\phi_A) \in \mathrm{Bimon}^R$, let $\mathbf{F}(A,\phi_A)$ be the functor indicated by Lemma 3; note in particular that $A = \mathbf{F}(A,\phi(A))(1)$ and $\phi(r) = \mathbf{F}(A,\phi_A)(r : 1 \to 1)$ for all $r \in R$. Suppose $f : (A,\phi_A) \to (B,\phi_B)$ is a morphism in Bimon^R. We must define a natural transformation $\mathbf{F}(f)$ from $\mathbf{F}(A,\phi_A)$ to $\mathbf{F}(B,\phi_B)$; let $\mathbf{F}(f)(n) = f^{\otimes n}$. We need to show that for every $n \times m$ matrix X with coefficients in R, the diagram

$$\begin{tikzcd}
A^{\otimes m} \arrow[r, "F(A)(X)"] \arrow[d, "f^{\otimes m}"'] & A^{\otimes n} \arrow[d, "f^{\otimes n}"] \\
B^{\otimes m} \arrow[r, "F(B)(X)"'] & B^{\otimes n}
\end{tikzcd}$$

commutes. Because of the enriched structure on the functors $\mathbf{F}(A,\phi_A)$ and $\mathbf{F}(B,\phi_B)$, it suffices to check this condition for matrices with precisely one nonzero entry. For 1×1 matrices (r), this becomes $f\phi_A(r) = \phi_B(r)(f)$, which is true since f is a morphism in Bimon^R. It is also true when m or n is 0 since $(\mathbf{F}(A,\phi_A))(0)$ is both an initial and terminal object. Since each matrix with one nonzero entry is a tensor product of matrices of these types, we know that $\mathbf{F}$ is a functor.

Now we want a functor $\mathbf{G}$ from the category of algebras over $\mathrm{Mat}(R)$ to Bimon^R; letting $\mathbf{G}$ be the functor indicated by Lemma 4, we have $\mathbf{G}(F : \mathrm{Mat}(R) \to \mathscr{C}) = (F(1), F(\mathrm{Hom}_{\mathrm{Mat}(R)}(1,1)))$. Our functor sends the natural transformation $\alpha : F \to G$ to the component α_1. This is well-defined since

$$\begin{tikzcd}
F(1) \arrow[r, "F(r)"] \arrow[d, "\alpha_1"'] & F(1) \arrow[d, "\alpha_1"] \\
G(1) \arrow[r, "G(r)"'] & G(1)
\end{tikzcd}$$

commutes. We know $\mathbf{G}$ is a functor because composition of natural transformations is defined componentwise.

By the way we have defined $\mathbf{F}$ and $\mathbf{G}$, we can see that $\mathbf{GF} = \mathrm{id}_{\mathrm{Bimon}^R}$ and $\mathbf{FG}$ is the identity functor on the category of algebras of $\mathrm{Mat}(R)$, so we have an isomorphism of categories. ☐

The fact that $\phi : R \to \mathrm{End}(A)$ is a rig homomorphism has the following diagrammatic representation, letting a triangle containing the variable $r \in R$ represent $\phi(r)$:



These diagrams, together with those in Section 2, constitute the entire set of relations given by Baez and Erbele for $\mathrm{Mat}(k)$, where k is a field [5].

5 Examples

As noted earlier, $\mathrm{Mat}(R)$ is equivalent to some famous PROPs for the right choices of R. The only possible two-element rigs are the Boolean rig $\mathbb{B}$ where $1+1=1$ and $\mathbb{F}_2$ where $1+1=0$. We define FinRel to be the category whose objects are finite sets and whose morphisms are relations between them. Because all sets of the same cardinality are isomorphic in this category, we can make FinRel into a PROP by choosing one representative of each cardinality and defining the tensor product on objects to be the disjoint union of sets. A relation from m to n can be defined as a Boolean-valued matrix by letting the ith row and jth column be 1 if and only if i relates to j. Relations can also be visualized as string diagrams, where i relates to j if and only if a path connects the two; an example is given in Figure 1.



Figure 1: A string diagram for a relation

If A is an algebra over $\mathrm{Mat}(\mathbb{B})$ equippped with a map of rigs $\phi : \mathbb{B} \to \mathrm{End}(A)$ and $a \in \mathrm{End}(A)$, then $a+a = (\phi(1_R)+\phi(1_R))a = a$; conversely, if $a+a=a$ for all $a \in \mathrm{End}(A)$ then $\phi(1_R)+\phi(1_R)=\phi(1_R)$, so the algebras of $\mathrm{Mat}(\mathbb{B})$ are exactly those where $a+a=a$ for all $a \in \mathrm{End}(A)$.

Let A be a bimonoid; we say A is *special* if $\mu_A\Delta_A = \mathrm{id}_A$. Note that

$$a+a = \mu_A(a\otimes a)\Delta_A = \mu_A\Delta_A a,$$

so it follows that $a+a=a$ for all $a \in \mathrm{End}(A)$ if and only if A is special. Therefore we have

Corollary 6. *The category* FinRel *whose objects are finite sets and whose morphisms are relations between them is the PROP for special bicommutative bimonoids.*

Replacing $1_R+1_R=1_R$ with $1_R+1_R=0_R$ gives us $a+a=0$ for all $a\in\mathrm{End}(A)$, so the above discussion yields:

Corollary 7. Mat$(\mathbb{F}_2)$ *is the PROP for bicommutative bimonoids A where* $\mu_A\Delta_A = \eta_A\varepsilon_A$.

Now consider the case Mat$(\mathbb{N})$. This category is equivalent to FinSpan whose objects are finite sets and whose morphisms $X \to Y$ are triples $(S, f : S \to X, g : S \to Y)$, where S is a set and f, g are ordinary functions. To see this, if M is a $\mathbb{N}$-valued matrix, we can think of m_{ij} as the cardinality of the set

$$\{s \in S | f(s) = j \text{ and } g(s) = i\}.$$

Since $\mathbb{N}$ is an inital object in the category of rigs,

Corollary 8. FinSpan *is the PROP for bicommutative bimonoids.*

An equivalent version of this result was proved by Lack [4, 9].

For a symmetric monoidal category $\mathscr{C}$, define a *Hopf monoid* A over $\mathscr{C}$ to be a bimonoid equipped with an additional morphism $S : A \to A$ satisfying

$$\mu_A(S \otimes \mathrm{id}_A)\Delta_A = \eta_A\varepsilon_A = \mu_A(\mathrm{id}_A \otimes S)\Delta_A.$$

In the case where $\mathscr{C}$ is the category of vector spaces over a field k, we call A a *Hopf algebra*; thorough explainations of Hopf algebras are given in [6, 13]. We know that S is a bimodule antihomomorphism [13], so if A is bicommutative, S is a bimodule homomorphism. Therefore the enriched structure on End(A) simplifies the above axiom to

$$S + \mathrm{id}_A = 0_A.$$

Now that we know that S is a bimodule homomorphism, we can say that

$$0 = \eta_A\varepsilon_A = S\eta_A\varepsilon_A = S\mu_A(S \otimes \mathrm{id}_A)\Delta_A = \mu(S^2 \otimes S)\Delta_A = S^2 + S,$$

from which it follows that $S^2 = \mathrm{id}_A$.

Corollary 9. Mat$(\mathbb{Z})$ *is the PROP for bicommutative Hopf monoids.*

Proof. We must show that a bicommutative Hopf monoid is exactly a bicommutative bialgebra A equipped with a rig homomorphism $\phi : R \to \mathrm{End}(A)$.

If A is such an algebra, then A is a bicommutative Hopf monoid with antipode $\phi(-1)$, since

$$\eta_A\varepsilon_A = \phi(0) = \phi(1) + \phi(-1) = \mu_A(\mathrm{id}_A \otimes \phi(-1))\Delta_A$$

and

$$\eta_A\varepsilon_A = \phi(0) = \phi(-1) + \phi(1) = \mu_A(\phi(-1) \otimes \mathrm{id}_A)\Delta_A.$$

Conversely, Let A be a bicommutative Hopf monoid with antipode S. We need to show that $\phi(-1) = S$ extends uniquely to a rig homomorphism $\phi : \mathbb{Z} \to \mathrm{End}(A)$. We are locked into choosing $\phi(0) = 0$, $\phi(n) = \mathrm{id}_A + \phi(n-1)$, and $\phi(-n) = S + \phi(-n+1)$ inductively for each positive integer n. Since $S + \mathrm{id}_A = 0$ is the antipode axiom, we have that ϕ is an additive homomorphism of commutative monoids. That ϕ is a homomorphism of rigs now follows from the distributive law for rigs and the fact that $S^2 = \mathrm{id}_A$ since A is a bicommutative Hopf monoid. ☐

Acknowledgements

We thank John Baez for his help and for his “Theorems into Coffee” series of blog posts, which posed this paper’s central questions [1, 2, 3, 4]. We also thank Jason Erbele for allowing us to use his diagrams and for his assistance with TikZ.

References

[1] J. Baez, Theorems into Coffee, weblog post at *The n-Category Café* (May 4, 2008). Available at https://golem.ph.utexas.edu/category/2008/05/theorems_into_coffee.html.

[2] J. Baez, Theorems into Coffee II, weblog post at *The n-Category Café* (May 6, 2008). Available at https://golem.ph.utexas.edu/category/2008/05/theorems_into_coffee_ii.html.

[3] J. Baez, Theorems into Coffee III, weblog post at *The n-Category Café* (May 15, 2008). Available at https://golem.ph.utexas.edu/category/2008/05/theorems_into_coffee_iii.html.

[4] J. Baez, Theorems into Coffee IV, weblog post at *The n-Category Café* (July 12, 2008). Available at https://golem.ph.utexas.edu/category/2008/07/theorems_into_coffee_iv.html.

[5] J. Baez and J. Erbele, Categories in control. Available as arXiv:1405.6881.

[6] G. Bergman, Everybody knows what a Hopf algebra is, *Amer. Math. Soc. Contemp. Math.* **43** (1985), 25–48.

[7] F. Bonchi, P. Sobociński and F. Zanasi, Interacting Hopf algebras. Available as arXiv:1403.7048.

[8] F. Bonchi, P. Sobociński and F. Zanasi, A categorical semantics of signal flow graphs, in *CONCUR 2014–Concurrency Theory*, eds. P. Baldan and D. Gorla, *Lecture Notes in Computer Science* vol. 8704, Springer, Berlin, 2014, pp. 435–450. Also available at http://users.ecs.soton.ac.uk/ps/papers/sfg.pdf.

[9] S. Lack, Composing PROPs, *Th. Appl. Cat.* **13** (2004), 147–163. Available at http://www.tac.mta.ca/tac/volumes/13/9/13-09.pdf.

[10] S. Mac Lane, Categorical algebra, *Bull. Amer. Math. Soc.* **71** (1965), 40–106. Available at http://www.ams.org/journals/bull/1965-71-01/S0002-9904-1965-11234-4/S0002-9904-1965-11234-4.pdf.

[11] M. Markl, Algebras and PROPs. Available as arXiv:0601129.

[12] T. Pirashvili, On the PROP corresponding to bialgebras, *Cahiers Top. Géom. Diff. Cat.* **43** (2002), 221–239. Available at http://www.numdam.org/numdam-bin/item?id=CTGDC_2002__43_3_221_0.

[13] M. Sweedler, *Hopf Algebras*, W. A. Benjamin, Inc., 1969.